
Standing Orders for Administering Hepatitis B Vaccine to Adults

Purpose: To reduce morbidity and mortality from hepatitis B virus (HBV) infection by vaccinating all adults who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices.

Policy: Under these standing orders, eligible nurses and other healthcare professionals (e.g., pharmacists), where allowed by state law, may vaccinate adults who meet any of the criteria below.

Procedure:

1. Identify adults in need of hepatitis B vaccination based on the following criteria:
 - a. Age younger than 19 years with no or unknown history of prior receipt of a complete series of hepatitis B vaccine
 - b. Age 19 years or older meeting any of the following criteria:
 - patient with end-stage renal disease, including patients receiving hemodialysis
 - patient with HIV infection
 - patient with chronic liver disease
 - sexually active and not in a long-term, mutually monogamous relation (i.e., more than 1 sex partner during the previous 6 months)
 - under evaluation or treatment for a sexually transmitted disease (STD)
 - a male who has sex with males
 - current or recent injection-drug user
 - at occupational risk of infection through exposure to blood or blood-contaminated body fluids (e.g., healthcare worker, public safety worker, trainee in a health professional or allied health school)
 - client or staff of an institution for persons with developmental disabilities
 - sex partner or household member of a person who is chronically infected with HBV (including an HBsAg-positive adopted child)
 - planned travel to a country with high or intermediate prevalence of chronic HBV infection (a list of countries is available at www.cdc.gov/travel/diseases.htm)
 - housed in or seen for care in a setting in which a high proportion of persons have risk factors for HBV infection (e.g., STD treatment facilities, correctional facilities, institutions for developmentally disabled persons)
 - c. Any person who wishes to be vaccinated against HBV infection
2. Screen all patients for contraindications and precautions to hepatitis B vaccine:
 - a. **Contraindication:** a history of a serious reaction (e.g., anaphylaxis) after a previous dose of hepatitis B vaccine or to a hepatitis B vaccine component. For a list of vaccine components, go to www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf.
 - b. **Precaution:** moderate or severe acute illness with or without fever
3. Provide all patients with a copy of the most current federal Vaccine Information Statement (VIS). You must document, in the patient's medical record or office log, the publication date of the VIS and the date it was given to the patient. Provide non-English speakers with the VIS in their native language, if available; these can be found at www.immunize.org/vis.
4. Administer hepatitis B vaccine intramuscularly (22–25g, 1–1½" needle) in the deltoid muscle. For persons age 20 years or older, give 1.0 mL dosage; for persons age 19 years or younger, give 0.5 mL dosage.
5. Provide subsequent doses of hepatitis B vaccine to complete each patient's 3-dose schedule by observing a minimum interval of 4 weeks between the first and second doses, 8 weeks between the second and third doses, and at least 4 months (16 weeks) between the first and third doses.
6. Document each patient's vaccine administration information and follow up in the following places:
 - a. **Medical chart:** Record the date the vaccine was administered, the manufacturer and lot number, the vaccination site and route, and the name and title of the person administering the vaccine. If vaccine was not given, record the reason(s) for non-receipt of the vaccine (e.g., medical contraindication, patient refusal).
 - b. **Personal immunization record card:** Record the date of vaccination and the name/location of the administering clinic.
7. Be prepared for management of a medical emergency related to the administration of vaccine by having a written emergency medical protocol available, as well as equipment and medications.
8. Report all adverse reactions to hepatitis B vaccine to the federal Vaccine Adverse Event Reporting System (VAERS) at www.vaers.hhs.gov or by calling (800) 822-7967. VAERS report forms are available at www.vaers.hhs.gov.

*For persons born in Asia, the Pacific Islands, Africa, or other countries identified as having high rates of HBV infection (see MMWR 2005;54 [No. RR-16]:25), ensure that they have also been tested for hepatitis B surface antigen (HBsAg) to find out if they are chronically infected. If test is performed on same visit, give hepatitis B vaccine after the blood draw. Do not delay initiating hepatitis B vaccination while waiting for test results. If patient is found to be HBsAg-positive, appropriate medical follow-up should be provided.

This policy and procedure shall remain in effect for all patients of the _____ until rescinded or until _____ (date). (name of practice or clinic)

Medical Director's signature: _____ Effective date: _____