

# Standing Orders Templates for Administering Vaccines to Children/Teens and Adults

Visit [www.immunize.org/standing-orders](http://www.immunize.org/standing-orders) for all sets.

► Click blue text to view standing orders documents

Download these standing orders and use them “as is” or modify them to suit your work setting.

WEIGHT AND WEIGHT OF PATIENT	NEEDLE GAUGE	NEEDLE LENGTH
Female or male less than 130 lbs	22–25	5/8"–1"
Female or male 130–152 lbs	22–25	1"
Female 153–200 lbs	22–25	1–1½"
Male 153–260 lbs	22–25	1–1½"
Female 200+ lbs	22–25	1½"
Male 260+ lbs	22–25	1½"

25-gauge needle may be used in patients weighing less than 130 lbs (if the needle is held tight, the subcutaneous tissue is not penetrated).

Standing orders for other vaccines are available at [www.immunize.org/standing-orders](http://www.immunize.org/standing-orders). Note: This standing orders template may be adapted per a practice's discretion without obtaining permission from IAC. As a courtesy, please acknowledge IAC as its source.

## STANDING ORDERS FOR Administering Pneumococcal Vaccines (PCV13 and PPSV23) to Adults

### Purpose

To reduce morbidity and mortality from pneumococcal disease by vaccinating all adults who meet the criteria established by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices.

### Policy

Where allowed by state law, standing orders enable eligible nurses and other health care professionals (e.g., pharmacists) to assess the need for vaccination and to vaccinate adults who meet any of the criteria below.

### Procedure

**1 Assess Adults for Need of Vaccination** against *Streptococcus pneumoniae* (pneumococcus) infection according to the following criteria:

**Routine pneumococcal vaccination** – Assess adults age 65 years or older for need of pneumococcal vaccination. Pneumococcal conjugate vaccine (PCV13) should be administered routinely to all previously unvaccinated adults age 65 years and older. Pneumococcal polysaccharide vaccine (PPSV23) is recommended for all adults ages 65 years or older. For complete details, see section 5 (page 2).

**Risk-based pneumococcal vaccination** – Age 19 through 64 years with an underlying medical condition or other risk factor as described in the following table:

CATEGORY OF UNDERLYING MEDICAL CONDITION OR OTHER RISK FACTOR	RECOMMENDED VACCINES ARE MARKED "X" BELOW		
	PCV13	PPSV23	PPSV23 booster <sup>a</sup>
Chronic heart disease, <sup>1</sup> chronic lung disease <sup>2</sup>		x	
Diabetes mellitus		x	
Chronic liver disease, cirrhosis		x	
Cigarette smoking		x	
Alcoholism		x	
Cochlear implant, cerebrospinal fluid leak	x	x	
Sickle cell disease, other hemoglobinopathy	x	x	x
Congenital or acquired asplenia	x	x	x
Congenital or acquired immunodeficiency, <sup>3</sup> HIV	x	x	x
Chronic renal failure, nephrotic syndrome	x	x	x
Leukemia, lymphoma	x	x	x
Generalized malignancy, Hodgkin disease	x	x	x
Idiopathic immunosuppression <sup>4</sup>	x	x	x
Solid organ transplant, multiple myeloma	x	x	x

<sup>a</sup> a second dose 5 years after the first dose of PPSV23

<sup>1</sup> Excluding hypertension  
<sup>2</sup> Including asthma  
<sup>3</sup> Including B (humoral) or T-lymphocyte deficiency, complement deficiencies (particularly C1, C2, C3, and C4 deficiencies), and phagocytic disorders (excluding chronic granulomatous disease)  
<sup>4</sup> Diseases requiring treatment with immunosuppressive drugs, including long term systemic corticosteroids and radiation therapy

### 2 Screen for Contraindications and Precautions

**Contraindications** – Do not give pneumococcal vaccine (PCV13 or PPSV23) to a person who has experienced a serious systemic or anaphylactic reaction to a prior dose of the vaccine or to any of its components. For a list of vaccine components, refer to the manufacturer's package insert ([www.immunize.org/packageinserts](http://www.immunize.org/packageinserts)) or go to [www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/exipient-table-2.pdf](http://www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/exipient-table-2.pdf).

**Precautions** – Moderate or severe acute illness with or without fever

CONTINUED ON THE NEXT PAGE ►

**IMMUNIZATION ACTION COALITION** Saint Paul, Minnesota • 651-647-9009 • [www.immunize.org](http://www.immunize.org) • [www.vaccineinformation.org](http://www.vaccineinformation.org)  
[www.immunize.org/catg.d/p1075.pdf](http://www.immunize.org/catg.d/p1075.pdf) • Item #P93075 (1/1/15)

STANDING ORDER (date of latest revision)	VACCINES	STANDING ORDER (date of latest revision)
<a href="#">child</a> (OCT 2012)	DTaP	—
<a href="#">child/teen</a> (JUNE 2013)	HepA	<a href="#">adult</a> (JUNE 2013)
<a href="#">child/teen</a> (OCT 2012)	HepB	<a href="#">adult</a> (OCT 2015)
<a href="#">child</a> (JUNE 2015)	Hib	<a href="#">adult</a> (JUNE 2015)
<a href="#">child/teen</a> (JAN 2017)	HPV	<a href="#">adult</a> (JAN 2017)
<a href="#">child/teen</a> (OCT 2014)	IPV (polio)	—
<a href="#">child/teen</a> (SEPT 2016)	Influenza	<a href="#">adult</a> (SEPT 2016)
<a href="#">child/teen</a> (JUNE 2013)	MMR	<a href="#">adult</a> (JUNE 2013)
<a href="#">child/teen</a> (DEC 2016)	MenACWY (MCV4), MPSV	<a href="#">adult</a> (DEC 2016)
<a href="#">teen</a> (NOV 2016)	MenB	<a href="#">adult</a> (NOV 2016)
<a href="#">child/teen</a> (APRIL 2013)	PCV	<a href="#">adult</a> (JAN 2017)
<a href="#">child</a> (MAY 2015)	PPSV	
<a href="#">child</a> (FEB 2014)	Rotavirus	—
—	Tdap	<a href="#">pregnant woman</a> (FEB 2014)
<a href="#">child/teen</a> (APRIL 2013)	Tdap/Td	<a href="#">adult</a> (JAN 2017)
<a href="#">child/teen</a> (JULY 2016)	Varicella	<a href="#">adult</a> (JULY 2016)
—	Zoster	<a href="#">adult</a> (NOV 2015)

All sets of standing orders for routinely recommended vaccines are available at [www.immunize.org/standing-orders](http://www.immunize.org/standing-orders)