


Figure 3. Vaccines that might be indicated for children and adolescents ages 18 years or younger based on medical indications

Vaccine ▼	Indication ►	Pregnancy	Immunocompromised status (excluding HIV infection)	HIV infection CD4+ count (cells/ μ L)		Kidney failure, end-stage renal disease, on hemodialysis	Heart disease, chronic lung disease	CSF leaks/cochlear implants	Asplenia and persistent complement component deficiencies	Chronic liver disease	Diabetes
				<15% of total CD4 cell count	\geq 15% of total CD4 cell count						
Hepatitis B ¹											
Rotavirus ²			SCID*								
Diphtheria, tetanus, and acellular pertussis ³ (DTaP)											
<i>Haemophilus influenzae</i> type b ⁴											
Pneumococcal conjugate ⁵											
Inactivated poliovirus ⁶											
Influenza ⁷											
Measles, mumps, rubella ⁸											
Varicella ⁹											
Hepatitis A ¹⁰											
Meningococcal ACWY ¹¹											
Tetanus, diphtheria, and acellular pertussis ¹² (Tdap)											
Human papillomavirus ¹³											
Meningococcal B ¹¹											
Pneumococcal polysaccharide ⁵											



*Severe Combined Immunodeficiency

Footnotes: Recommended Immunization Schedule for Children and Adolescents Ages 18 Years or Younger, United States, 2017

For further guidance on the use of the vaccines mentioned below, see www.cdc.gov/vaccines/hcp/acip-recs/index.html.

For vaccine recommendations for persons ages 19 years and older, see the Recommended Adult Immunization Schedule.

1. Hepatitis B (HepB) vaccine. (Minimum age: birth)

Routine vaccination:

- At birth:
 - Administer monovalent HepB vaccine to all newborns within 24 hours of birth.
 - For infants born to hepatitis B surface antigen (HBsAg)-positive mothers, administer HepB vaccine and 0.5 mL of hepatitis B immune globulin (HBIG) within 12 hours of birth. These infants should be tested for HBsAg and antibody to HBsAg (anti-HBs) at age 9 through 12 months (preferably at the next well-child visit) or 1 to 2 months after completion of the HepB series if the series was delayed.
 - If mother's HBsAg status is unknown, within 12 hours of birth, administer HepB vaccine regardless of birth weight. For infants weighing less than 2,000 grams, administer HBIG in addition to HepB vaccine within 12 hours of birth. Determine mother's HBsAg status as soon as possible and, if mother is HBsAg positive, also administer HBIG to infants weighing 2,000 grams or more as soon as possible, but no later than age 7 days.

Doses following the birth dose:

- The second dose should be administered at age 1 or 2 months. Monovalent HepB vaccine should be used for doses administered before age 6 weeks.
- Infants who did not receive a birth dose should receive 3 doses of a HepB-containing vaccine on a schedule of 0, 1 to 2 months, and 6 months, starting as soon as feasible (see Figure 2).
- Administer the second dose 1 to 2 months after the first dose (minimum interval of 4 weeks); administer the third dose at least 8 weeks after the second dose and at least 16 weeks after the first dose. The final (third or fourth) dose in the HepB vaccine series should be administered **no earlier than age 24 weeks**.
- Administration of a total of 4 doses of HepB vaccine is permitted when a combination vaccine containing HepB is administered after the birth dose.

Catch-up vaccination:

- Unvaccinated persons should complete a 3-dose series.
- A 2-dose series (doses separated by at least 4 months) of adult formulation Recombivax HB is licensed for use in children age 11 through 15 years.
- For other catch-up guidance, see Figure 2.

2. Rotavirus (RV) vaccines. (Minimum age: 6 weeks for both RV1 [Rotarix] and RV5 [RotaTeq])

Routine vaccination:

- Administer a series of RV vaccine to all infants as follows:
 - If Rotarix is used, administer a 2-dose series at ages 2 and 4 months.
 - If RotaTeq is used, administer a 3-dose series at ages 2, 4, and 6 months.
 - If any dose in the series was RotaTeq or vaccine product is unknown for any dose in the series, a total of 3 doses of RV vaccine should be administered.

Catch-up vaccination:

- The maximum age for the first dose in the series is 14 weeks, 6 days; vaccination should not be initiated for infants ages 15 weeks, 0 days, or older.
- The maximum age for the final dose in the series is 8 months, 0 days.
- For other catch-up guidance, see Figure 2.

3. Diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine. (Minimum age: 6 weeks. Exception: DTaP-IPV [Kinrix, Quadracel]: 4 years)

Routine vaccination:

- Administer a 5-dose series of DTaP vaccine at ages 2, 4, 6, 15 through 18 months, and 4 through 6 years. The fourth dose may be administered as early as age 12 months, provided at least 6 months have elapsed since the third dose.
- Inadvertent administration of fourth DTaP dose early: If the fourth dose of DTaP was administered at least 4 months after the third dose of DTaP and the child was age 12 months or older, it does not need to be repeated.

Catch-up vaccination:

- The fifth dose of DTaP vaccine is not necessary if the fourth dose was administered at age 4 years or older.
- For other catch-up guidance, see Figure 2.

4. Haemophilus influenzae type b (Hib) conjugate vaccine. (Minimum age: 6 weeks for PRP-T [ActHIB, DTaP-IPV/Hib (Pentacel), Hiberix, and Hib-MenCY (MenHibrix)], PRP-OMP [PedvaxHIB])

Routine vaccination:

- Administer a 2- or 3-dose Hib vaccine primary series and a booster dose (dose 3 or 4, depending on vaccine used in primary series) at age 12 through 15 months to complete a full Hib vaccine series.

(continued)