

Summary of Recommendations for Adult Immunization

Vaccine name and route	For whom vaccination is recommended	Schedule for vaccine administration (any vaccine can be given with another)	Contraindications and precautions (mild illness is not a contraindication)
<p>Seasonal Influenza Trivalent inactivated influenza vaccine (TIV) <i>Give IM</i></p> <p>Live attenuated influenza vaccine (LAIV) <i>Give intranasally</i></p>	<ul style="list-style-type: none"> Vaccinate all persons who want to reduce the risk of becoming ill with influenza or spreading it to others. Special efforts should be made to vaccinate the following persons because they are at higher risk for influenza complications: those who are ages 50yrs and older; have pulmonary (including asthma), cardiovascular (except hypertension), renal, hepatic, cognitive, neurologic/neuromuscular, hematologic, or metabolic (including diabetes) disorders; have immunosuppression (including that caused by medications or HIV); will be pregnant during influenza season; residing in long-term care facilities. Vaccinate adults, children, and teens who are household contacts, caregivers, or workplace contacts of the persons listed in bullet #2 above or of children age 0–59m. Vaccinate healthcare personnel. Travelers to the tropics, to areas with current influenza activity, or on trips with people from areas with current influenza activity (e.g., on organized tours) should consider vaccination. Vaccinate students or other persons in institutional settings (e.g., dormitories, correctional facilities). <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> <p>Note: LAIV may not be given to some of the persons listed to the left; see contraindications listed in far right column.</p> </div>	<ul style="list-style-type: none"> Give 1 dose every year in the fall or winter. Begin vaccination services as soon as vaccine is available and continue until the supply is depleted. Continue to give vaccine to unvaccinated adults throughout the influenza season (including when influenza activity is present in the community) and at other times when the risk of influenza exists. If 2 or more of the following live virus vaccines are to be given—LAIV, MMR, Var, Zos, and/or yellow fever vaccine—they should be given on the same day. If they are not, space them by at least 28d. 	<p>Contraindications</p> <ul style="list-style-type: none"> Previous anaphylactic reaction to this vaccine, to any of its components, or to eggs. For LAIV only: age 50yrs and older; pregnancy; chronic pulmonary (including asthma), cardiovascular (except hypertension), renal, hepatic, neurological/neuromuscular, hematologic, or metabolic (including diabetes) disorders; immunosuppression (including that caused by medications or HIV). <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. History of Guillain-Barré syndrome (GBS) within 6wks of previous influenza vaccination. For LAIV only: close contact with an immunosuppressed person when the person requires protective isolation.
<p>Pneumococcal polysaccharide (PPSV) <i>Give IM or SC</i></p>	<ul style="list-style-type: none"> Persons age 65yrs and older. Persons who have chronic illness or other risk factors, including chronic cardiac or pulmonary disease (including asthma), chronic liver disease, alcoholism, diabetes, CSF leaks, cigarette smoking, as well as people living in special environments or social settings (including Alaska Natives and certain American Indian populations age 50 through 64yrs if recommended by local public health authorities). Those at highest risk of fatal pneumococcal infection, including persons who <ul style="list-style-type: none"> Have anatomic asplenia, functional asplenia, or sickle cell disease. Have an immunocompromising condition, including HIV infection, leukemia, lymphoma, Hodgkin’s disease, multiple myeloma, generalized malignancy, chronic renal failure, or nephrotic syndrome. Are receiving immunosuppressive chemotherapy (including corticosteroids). Have received an organ or bone marrow transplant. Are candidates for or recipients of cochlear implants. 	<ul style="list-style-type: none"> Give 1 dose if unvaccinated or if previous vaccination history is unknown. Give a 1-time revaccination at least 5yrs after 1st dose to persons <ul style="list-style-type: none"> Age 65yrs and older if the 1st dose was given prior to age 65yrs At highest risk of fatal pneumococcal infection or rapid antibody loss (see the 3rd bullet in the box to left for listings of persons at highest risk). 	<p>Contraindication</p> <p>Previous anaphylactic reaction to this vaccine or to any of its components.</p> <p>Precaution</p> <p>Moderate or severe acute illness.</p>
<p>Zoster (shingles) (Zos) <i>Give SC</i></p>	<ul style="list-style-type: none"> Persons age 60yrs and older. 	<ul style="list-style-type: none"> Give 1-time dose if unvaccinated, regardless of previous history of herpes zoster (shingles) or chickenpox. If 2 or more of the following live virus vaccines are to be given—MMR, Zos, and/or yellow fever vaccine—they should be given on the same day. If they are not, space them by at least 28d. 	<p>Contraindications</p> <ul style="list-style-type: none"> Previous anaphylactic reaction to any component of zoster vaccine (e.g., gelatin & neomycin). Primary cellular or acquired immunodeficiency. Pregnancy. <p>Precaution</p> <p>Moderate or severe acute illness.</p>

*This document was adapted from the recommendations of the Advisory Committee on Immunization Practices (ACIP). To obtain copies of these recommendations, call the CDC-INFO Contact Center at (800) 232-4636; visit CDC’s website at www.cdc.gov/vaccines/pubs/ACIP-list.htm; or visit the Immunization Action Coal-

ition (IAC) website at www.immunize.org/acip. This table is revised periodically. Visit IAC’s website at www.immunize.org/adultrules to make sure you have the most current version.

Summary of Recommendations for Adult Immunization (continued)

Vaccine name and route	For whom vaccination is recommended	Schedule for vaccine administration (any vaccine can be given with another)	Contraindications and precautions (mild illness is not a contraindication)
<p>Hepatitis B (HepB) <i>Give IM</i></p> <p>Brands may be used interchangeably.</p>	<ul style="list-style-type: none"> All persons through age 18yrs. All adults wishing to be protected from hepatitis B virus infection. High-risk persons, including household contacts and sex partners of HBsAg-positive persons; injecting drug users; sexually active persons not in a long-term, mutually monogamous relationship; men who have sex with men; persons with HIV; persons seeking evaluation or treatment for an STD; patients receiving hemodialysis and patients with renal disease that may result in dialysis; healthcare personnel and public safety workers who are exposed to blood; clients and staff of institutions for the developmentally disabled; inmates of long-term correctional facilities; and certain international travelers. Persons with chronic liver disease. <p>Note: Provide serologic screening for immigrants from endemic areas. If patient is chronically infected, assure appropriate disease management. Screen sex partners and household members; give HepB at the same visit if not already vaccinated.</p>	<ul style="list-style-type: none"> Give 3 doses on a 0, 1, 6m schedule. Alternative timing options for vaccination include 0, 2, 4m and 0, 1, 4m. There must be at least 4wks between doses #1 and #2, and at least 8wks between doses #2 and #3. Overall, there must be at least 16wks between doses #1 and #3. Schedule for those who have fallen behind: If the series is delayed between doses, DO NOT start the series over. Continue from where you left off. <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> <p>For Twinrix (hepatitis A and B combination vaccine [GSK]) for patients age 18yrs and older only: give 3 doses on a 0, 1, 6m schedule. There must be at least 4wks between doses #1 and #2, and at least 5m between doses #2 and #3. An alternative schedule can also be used at 0, 7d, 21–30d, and a booster at 12m.</p> </div>	<p>Contraindication Previous anaphylactic reaction to this vaccine or to any of its components.</p> <p>Precaution Moderate or severe acute illness.</p>
<p>Hepatitis A (HepA) <i>Give IM</i></p> <p>Brands may be used interchangeably.</p>	<ul style="list-style-type: none"> All persons wishing to be protected from hepatitis A virus (HAV) infection. Persons who travel or work anywhere EXCEPT the U.S., Western Europe, New Zealand, Australia, Canada, and Japan. Persons with chronic liver disease; injecting and non-injecting drug users; men who have sex with men; people who receive clotting-factor concentrates; persons who work with HAV in experimental lab settings; food handlers when health authorities or private employers determine vaccination to be appropriate. Persons who anticipate close personal contact with an international adoptee from a country of high or intermediate endemicity during the first 60 days following the adoptee's arrival in the U.S. Adults age 40yrs or younger with recent (within 2 wks) exposure to HAV. For persons older than age 40yrs with recent (within 2 wks) exposure to HAV, immune globulin is preferred over HepA vaccine. 	<ul style="list-style-type: none"> Give 2 doses. The minimum interval between doses #1 and #2 is 6m. If dose #2 is delayed, do not repeat dose #1. Just give dose #2. 	<p>Contraindication Previous anaphylactic reaction to this vaccine or to any of its components.</p> <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. Safety during pregnancy has not been determined, so benefits must be weighed against potential risk.
<p>Td, Tdap (Tetanus, diphtheria, pertussis) <i>Give IM</i></p>	<ul style="list-style-type: none"> All adults who lack written documentation of a primary series consisting of at least 3 doses of tetanus- and diphtheria-toxoid-containing vaccine. A booster dose of tetanus- and diphtheria-toxoid-containing vaccine may be needed for wound management as early as 5yrs after receiving a previous dose, so consult ACIP recommendations.* Using tetanus toxoid (TT) instead of Td or Tdap is <u>not</u> recommended. In pregnancy, when indicated, give Td or Tdap in 2nd or 3rd trimester. If not administered during pregnancy, give Tdap in immediate postpartum period. <p>For Tdap only:</p> <ul style="list-style-type: none"> All adults younger than age 65yrs who have not already received Tdap. Adults in contact with infants younger than age 12m (e.g., parents, grandparents younger than age 65yrs, childcare providers, healthcare personnel) who have not received a dose of Tdap should be prioritized for vaccination. Healthcare personnel who work in hospitals or ambulatory care settings and have direct patient contact and who have not received Tdap. 	<ul style="list-style-type: none"> For persons who are unvaccinated or behind, complete the primary series with Td (spaced at 0, 1–2m, 6–12m intervals). One-time dose of Tdap may be used for any dose if younger than age 65yrs. Give Td booster every 10yrs after the primary series has been completed. For adults younger than age 65yrs, a 1-time dose of Tdap is recommended to replace the next Td. Intervals of 2yrs or less between Td and Tdap may be used. <p>Note: The two Tdap products are licensed for different age groups: Adacel (sanofi) for use in persons age 11–64yrs and Boostrix (GSK) for use in persons age 10–64yrs.</p>	<p>Contraindications</p> <ul style="list-style-type: none"> Previous anaphylactic reaction to this vaccine or to any of its components. For Tdap only, history of encephalopathy within 7d following DTP/DTaP. <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. GBS within 6wks of receiving a previous dose of tetanus-toxoid-containing vaccine. Unstable neurologic condition. History of Arthus reaction following a previous dose of tetanus- and/or diphtheria-toxoid-containing vaccine, including MCV4. <p>Note: Tdap may be given to pregnant women at the provider's discretion.</p>
<p>Polio (IPV) <i>Give IM or SC</i></p>	<ul style="list-style-type: none"> Not routinely recommended for U.S. residents age 18yrs and older. <p>Note: Adults living in the U.S. who never received or completed a primary series of polio vaccine need not be vaccinated unless they intend to travel to areas where exposure to wild-type virus is likely. Previously vaccinated adults can receive 1 booster dose if traveling to polio endemic areas or to areas where the risk of exposure is high.</p>	<ul style="list-style-type: none"> Refer to ACIP recommendations* regarding unique situations, schedules, and dosing information. 	<p>Contraindication Previous anaphylactic or neurologic reaction to this vaccine or to any of its components.</p> <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. Pregnancy.

Summary of Recommendations for Adult Immunization (continued)

Vaccine name and route	For whom vaccination is recommended	Schedule for vaccine administration (any vaccine can be given with another)	Contraindications and precautions (mild illness is not a contraindication)
<p>Varicella (Var) (Chickenpox) <i>Give SC</i></p>	<ul style="list-style-type: none"> All adults without evidence of immunity. <p>Note: Evidence of immunity is defined as written documentation of 2 doses of varicella vaccine; a history of varicella disease or herpes zoster (shingles) based on healthcare-provider diagnosis; laboratory evidence of immunity; and/or birth in the U.S. before 1980, with the exceptions that follow. Healthcare personnel (HCP) and pregnant women born in the U.S. before 1980 who do not meet any of the criteria above should be tested. If they are not immune, give the 1st dose of varicella vaccine immediately (HCP) or postpartum and before hospital discharge (pregnant women). Give the 2nd dose 4–8 wks later. Routine post-vaccination serologic testing is not recommended.</p>	<ul style="list-style-type: none"> Give 2 doses. Dose #2 is given 4–8wks after dose #1. If dose #2 is delayed, do not repeat dose #1. Just give dose #2. If 2 or more of the following live virus vaccines are to be given—LAIV, MMR, Var, Zos, and/or yellow fever vaccine—they should be given on the same day. If they are not, space them by at least 28d. May use as postexposure prophylaxis if given within 5d. 	<p>Contraindications</p> <ul style="list-style-type: none"> Previous anaphylactic reaction to this vaccine or to any of its components. Pregnancy or possibility of pregnancy within 4wks. Persons on high-dose immunosuppressive therapy or who are immunocompromised because of malignancy and primary or acquired cellular immunodeficiency, including HIV/AIDS (although vaccination may be considered if CD4+ T-lymphocyte counts are greater than or equal to 200 cells/μL. See <i>MMWR 2007;56,RR-4</i>). <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. If blood, plasma, and/or immune globulin (IG or VZIG) were given in past 11m, see ACIP statement <i>General Recommendations on Immunization*</i> regarding time to wait before vaccinating.
<p>Meningococcal Conjugate vaccine (MCV4) <i>Give IM</i> Polysaccharide vaccine (MPSV4) <i>Give SC</i></p>	<ul style="list-style-type: none"> All persons age 11 through 18yrs. College freshmen living in a dormitory. Persons with anatomic or functional asplenia or with a persistent complement component deficiency. Persons who travel to or reside in countries in which meningococcal disease is hyperendemic or epidemic (e.g., the “meningitis belt” of Sub-Saharan Africa). Microbiologists routinely exposed to isolates of <i>N. meningitidis</i>. 	<ul style="list-style-type: none"> Give 1 dose. If previous vaccine was MCV4 or MPSV4, revaccinate after 5yrs if risk continues. MCV4 is preferred over MPSV4 for persons age 5yrs and younger; use MPSV4 ONLY if there is a permanent contraindication/precaution to MCV4. If the only risk factor is living in a campus dormitory, there is no need to give a 2nd dose. 	<p>Contraindication</p> <p>Previous anaphylactic or neurologic reaction to this vaccine or to any of its components, including diphtheria toxoid (for MCV4).</p> <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. For MCV4 only, history of Guillain-Barré syndrome (GBS).
<p>MMR (Measles, mumps, rubella) <i>Give SC</i></p>	<ul style="list-style-type: none"> Persons born in 1957 or later (especially those born outside the U.S.) should receive at least 1 dose of MMR if there is no laboratory evidence of immunity or documentation of a dose given on or after the first birthday. Persons in high-risk groups, such as healthcare personnel (paid, unpaid, or volunteer), students entering college and other post–high school educational institutions, and international travelers, should receive a total of 2 doses. Persons born before 1957 are usually considered immune, but evidence of immunity (serology or history of 2 doses of MMR) should be considered for healthcare personnel. Women of childbearing age who do not have acceptable evidence of rubella immunity or vaccination. <p>Note: Routine post-vaccination serologic testing is not recommended.</p>	<ul style="list-style-type: none"> Give 1 or 2 doses (see criteria in 1st and 2nd bullets in box to left). If dose #2 is recommended, give it no sooner than 4wks after dose #1. If a pregnant woman is found to be rubella susceptible, give 1 dose of MMR postpartum. If 2 or more of the following live virus vaccines are to be given—LAIV, MMR, Var, Zos, and/or yellow fever vaccine—they should be given on the same day. If they are not, space them by at least 28d. Within 72hrs of measles exposure, give 1 dose as postexposure prophylaxis to susceptible adults. 	<p>Contraindications</p> <ul style="list-style-type: none"> Previous anaphylactic reaction to this vaccine or to any of its components. Pregnancy or possibility of pregnancy within 4wks. Severe immunodeficiency (e.g., hematologic and solid tumors; receiving chemotherapy; congenital immunodeficiency; long-term immunosuppressive therapy; or severely symptomatic HIV.) Note: HIV infection is NOT a contraindication to MMR for those who are not severely immunocompromised (i.e., CD4+ T-lymphocyte counts are greater than or equal to 200 cells/μL). <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. If blood, plasma, and/or immune globulin were given in past 11m, see ACIP statement <i>General Recommendations on Immunization*</i> regarding time to wait before vaccinating. History of thrombocytopenia or thrombocytopenic purpura. <p>Note: If TST (tuberculosis skin test) and MMR are both needed but not given on same day, delay TST for 4–6wks after MMR.</p>
<p>Human papillomavirus (HPV) <i>Give IM</i></p>	<ul style="list-style-type: none"> All previously unvaccinated women through age 26yrs. 	<ul style="list-style-type: none"> Give 3 doses on a 0, 2, 6m schedule. There must be at least 4wks between doses #1 and #2 and at least 12wks between doses #2 and #3. Overall, there must be at least 24wks between doses #1 and #3. 	<p>Contraindication</p> <p>Previous anaphylactic reaction to this vaccine or to any of its components.</p> <p>Precautions</p> <ul style="list-style-type: none"> Moderate or severe acute illness. Data on vaccination in pregnancy are limited. Vaccination should be delayed until after completion of the pregnancy.