

Administering Vaccines: Dose, Route, Site, and Needle Size

Vaccine	Dose		Route
COVID-19 For product and dosage information for COVID-19 vaccine, see Immunize.org's "Checklist of Current Versions of U.S. COVID-19 Vaccination Guidance and Clinic Support Tools" at www.immunize.org/catg.d/p3130.pdf .			IM
Dengue (DENV4CYD)	0.5 mL		Subcut
Diphtheria, Tetanus, Pertussis (DTaP, Tdap, Td)	0.5 mL		IM
<i>Haemophilus influenzae</i> type b (Hib)	0.5 mL		IM
Hepatitis A (HepA)	≤18 yrs: 0.5 mL ≥19 yrs: 1.0 mL		IM
Hepatitis B (HepB) <i>People 11–15 yrs may be given Recombivax HB (Merck) 1.0 mL adult formulation on a 2-dose schedule.</i>	Engerix-B; Recombivax HB ≤19 yrs: 0.5 mL ≥20 yrs: 1.0 mL Heplisav-B, ≥18 yrs: 0.5 mL		IM
Human papillomavirus (HPV)	0.5 mL		IM
Influenza, live attenuated (LAIV)	0.2 mL (0.1 mL in each nostril)		Intranasal spray
Influenza, inactivated (IIV); 6 thru 35 mos • Egg-based IIV: Afluria, Fluzone, Fluarix, FluLaval • Cell-culture based (ccIIV): Flucelvax	Afluria: 0.25 mL Fluzone: 0.25 or 0.5 mL Fluarix, Flucelvax, FluLaval: 0.5 mL		IM
Influenza, inactivated (IIV) and • Cell-culture based (ccIIV), 3+ yrs; • Recombinant (RIV, Flublok), 18+ yrs; • Adjuvanted (aIIV, Fludax) 65+ yrs ¹ • High-dose (HD-IIV, Fluzone High Dose) 65+ yrs ¹	0.5 mL		IM
Measles, Mumps, Rubella (MMR)	0.5 mL	MMR II (Merck) Priorix (GSK)	IM or Subcut Subcut
Meningococcal serogroups A, C, W, Y (MenACWY)	0.5 mL		IM
Meningococcal serogroup B (MenB)	0.5 mL		IM
Mpox (Jynneos)	0.5 mL		Subcut ²
Pneumococcal conjugate (PCV)	0.5 mL		IM
Pneumococcal polysaccharide (PPSV23)	0.5 mL		IM or Subcut
Polio, inactivated (IPV)	0.5 mL		IM or Subcut
Respiratory Syncytial Virus (RSV) vaccine	0.5 mL		IM
RSV preventive antibody (nirsevimab)	0.5 mL, 1 mL, or 2 mL based on weight and/or age		IM
Rotavirus (RV)	Rotarix: 1.0 mL Rotateq: 2.0 mL		Oral
Varicella (VAR)	0.5 mL		IM or Subcut
Zoster (RZV)	Shingrix: 0.5 ³ mL		IM
Combination Vaccines			
DTaP-HepB-IPV (Pediarix) DTaP-IPV/Hib (Pentacel) DTaP-IPV (Kinrix; Quadracel) DTaP-IPV-Hib-HepB (Vaxelis)	0.5 mL		IM
MenABCWY (Penbraya; Penmenvy)	0.5 mL		IM
MMRV (ProQuad)	0.5 mL		IM or Subcut
HepA-HepB (Twinrix)	1.0 mL		IM

1 HD-IIV or aIIV are also options for solid organ transplant recipients 18–64 yrs on an immunosuppressive medication regimen.

2 Administration of Jynneos subcut (0.5 mL) is preferred. If an adult patient desires intradermal administration (0.1 mL), it is permitted under FDA emergency use authorization (see www.fda.gov/media/160774/download).

3 The Shingrix (RZV) vial may contain more than 0.5 mL. Do not administer more than 0.5 mL.

Intramuscular (IM) injection

Injection Site and Needle Size

Subcutaneous (Subcut) injection

Use a 23–25 gauge needle. Choose the injection site that is appropriate to the person's age and body mass.

AGE	NEEDLE LENGTH	INJECTION SITE
Infants (younger than 12 mos)	5/8"	Fatty tissue over anterolateral thigh muscle
Children 12 mos or older, adolescents, and adults	5/8"	Fatty tissue over anterolateral thigh muscle or fatty tissue over triceps

Intramuscular (IM) injection

Use a 22–25 gauge needle. Choose the injection site and needle length that is appropriate to the person's age and body mass.

AGE	NEEDLE LENGTH	INJECTION SITE
Newborns (1st 28 days)	5/8" ⁴	Anterolateral thigh muscle
Infants (1 through 11 mos)	1"	Anterolateral thigh muscle
Toddlers (1 through 2 yrs)	1–1 1/4"	Anterolateral thigh muscle ⁶
	5/8"–1"	Deltoid muscle of arm
Children (3 through 10 yrs)	5/8"–1"	Deltoid muscle of arm ⁶
	1–1 1/4"	Anterolateral thigh muscle
Adolescents and teens (11 through 18 yrs)	5/8"–1"	Deltoid muscle of arm ⁶
	1–1 1/2"	Anterolateral thigh muscle

Biological sex and weight of patient 19 yrs or older	NEEDLE LENGTH	INJECTION SITE
Female or male <130 lbs	5/8"–1"	Deltoid muscle of arm
Female or male 130–152 lbs	1"	Deltoid muscle of arm
Female 153–200 lbs Male 153–260 lbs	1–1 1/2"	Deltoid muscle of arm
Female more than 200 lbs Male more than 260 lbs	1 1/2"	Deltoid muscle of arm
Female or male, any weight	1 ⁵ –1 1/2"	Anterolateral thigh muscle


4 If skin is stretched tightly and subcutaneous tissues are not bunched.

5 Alternate needle lengths may be used if the skin is stretched tightly and subcutaneous tissues are not bunched, as follows: a) a 5/8" needle in toddlers, children, and patients weighing less than 130 lbs (less than 60 kg) for IM injection in the deltoid muscle only, or b) a 1" needle for administration in the thigh muscle for adults of any weight.

6 Preferred site

NOTE: Always refer to the package insert (found at www.immunize.org/official-guidance/fda/pkg-inserts) included with each biologic for complete vaccine administration information. CDC's Advisory Committee on Immunization Practices (ACIP) recommendations for the particular vaccine should be reviewed as well. Access the ACIP recommendations at www.immunize.org/official-guidance/cdc/acip-recs/vaccines.

Intranasal (NAS) administration of Flumist (LAIV) vaccine



Subcutaneous (Subcut) injection

