

Immunizations for Babies

A Guide for Parents

These are the vaccinations your baby needs!

At birth ^{1,2}	HepB	RSV-mAb ¹ 0–7 mos													
2 months ^{1,2}	HepB 1–2 mos ²	+	DTaP	+	PCV	+	Hib	+	Polio	+	RV				
4 months ^{1,2}	HepB ³	+	DTaP	+	PCV	+	Hib	+	Polio	+	RV				
6 months ^{1,2}	HepB 6–18 mos ²	+	DTaP	+	PCV	+	Hib ⁴	+	Polio 6–18 mos ²	+	RV ⁵	+	Influenza ⁶	+	COVID ⁷
12 months and older ^{1,2}	MMR 12–15 mos ²	+	DTaP 15–18 mos ²	+	PCV 12–15 mos ²	+	Hib 12–15 mos ²	+	Chickenpox 12–15 mos ²	+	HepA ⁸ 12–23 mos ²	+	Influenza ⁶	+	COVID ⁷

- Check with your doctor or nurse to make sure your baby is receiving all vaccinations on schedule. Many times vaccines are combined to reduce the number of injections. Be sure you ask for a record card with the dates of your baby's vaccinations; bring this with you to every visit.

Here's a list of the diseases your baby will be protected against:

HepB: hepatitis B, a serious liver disease

DTaP: diphtheria, tetanus (lockjaw), and pertussis (whooping cough)

PCV: pneumococcal conjugate vaccine protects against a serious blood, lung, and brain infection

Hib: *Haemophilus influenzae* type b, a serious brain, throat, and blood infection

Polio: polio, a serious paralyzing disease

RSV: respiratory syncytial virus, a serious lung infection

RV: rotavirus infection, a serious diarrheal disease

Influenza: a serious lung infection

MMR: measles, mumps, and rubella

COVID-19: a potentially serious infectious disease

HepA: hepatitis A, a serious liver disease

Chickenpox: also called varicella

Notes to above chart:

1. Infants age birth through 7 months may need RSV preventive antibody (RSV-mAb) before or during their first RSV season if RSV vaccine was not given during pregnancy. Certain older infants age 8 through 19 months may need RSV-mAb before their second RSV season.
2. This is the age range in which this vaccine should be given.
3. Your baby may not need a dose of Hep B vaccine at age 4 months, depending on the vaccine used. Check with your doctor or nurse.
4. Your baby may not need a dose of Hib vaccine at age 6 months, depending on the vaccine used. Check with your doctor or nurse.
5. Your baby may not need a dose of RV vaccine at age 6 months, depending on the vaccine used. Check with your doctor or nurse.
6. All children age 6 months and older should be vaccinated against influenza in the fall or winter of each year.
7. You and your child's doctor or nurse should discuss seasonal COVID-19 each year to decide if your child should be vaccinated.
8. Your child will need 2 doses of HepA vaccine, given at least 6 months apart.

