

# Meningococcal ACWY Vaccine Recommendations by Age and Risk Factor

A separate vaccine is needed for protection against meningococcal serogroup B disease.

MenACWY = Menactra, MenQuadfi (Sanofi Pasteur); Menveo (GlaxoSmithKline)  
MenACWY-D = Menactra MenACWY-CRM = Menveo MenACWY-TT = MenQuadfi

## Routine Recommendations for Use of Meningococcal A,C,W,Y Vaccine (MenACWY)

For preteens age 11 through 12 years	Give dose #1 of 2-dose MenACWY series. (Dose #2 is recommended at age 16 years.)
For teens age 13 through 15 years	Give catch-up dose #1 of 2-dose MenACWY series. (Dose #2 will be due at age 16 years.)
For teens at age 16 years	Give dose #2 of MenACWY.
Catch-up for teens age 17 through 18 years	If dose #2 not given at age 16 years, give dose #2 of MenACWY as catch-up.
Catch-up for teens age 16 through 18 years	If no history of prior vaccination with MenACWY, give 1 dose of MenACWY.
Catch-up for adults age 19 through 21 years	If no history of vaccination with MenACWY since the 16th birthday, consider giving 1 dose of MenACWY.
For first year college students living in residence halls	If student has either no history of prior vaccination with MenACWY, received only 1 dose when younger than age 16 years, or most recent dose given after 16th birthday and more than 5 years have elapsed, give 1 dose.

## Risk-based Recommendations for Persons with Underlying Medical Conditions or Other Risk Factors

TARGETED GROUP BY AGE/OR RISK FACTOR	PRIMARY DOSE(S) <sup>1</sup>	BOOSTER DOSE(S) <sup>1</sup>
<b>Travelers to or residents of countries where meningococcal disease is hyperendemic or epidemic, people present during outbreaks caused by a vaccine serogroup,<sup>2</sup> and other people with prolonged increased risk for exposure (e.g., microbiologists routinely working with <i>Neisseria meningitidis</i>)</b>		
For age 2 through 6 months	Give 3 doses of Menveo, 8 weeks apart, and a 4th dose at age 12–18 months. If possible, vaccination should begin at age 2 months.	If primary vaccination is completed before the 7th birthday: give one booster dose 3 years after primary series, then every 5 years thereafter, as long as risk remains. <sup>4</sup>  If primary vaccination is completed at age 7 years or older: give a booster dose every 5 years thereafter, as long as risk remains.
For age 7 through 23 months who have not initiated a series of MenACWY	If age 7–8 months, initiate 2-dose series of Menveo <sup>3</sup> or, if age 9–23 months, give either Menveo or Menactra. <sup>4</sup> Separate the 2 doses by at least 12 weeks. <sup>5</sup>	
For age 2 years and older	Give 1 dose of any MenACWY vaccine. <sup>4</sup>	
<b>People with persistent complement component deficiencies<sup>6</sup></b>		
For age 2 through 6 months	Give 3 doses of Menveo, 8 weeks apart, and a 4th dose at age 12–18 months. If possible, vaccination should begin at age 2 months.	If primary vaccination is completed before the 7th birthday: give one booster dose 3 years after primary series, then every 5 years thereafter, as long as risk remains. <sup>4</sup>  If primary vaccination is completed at age 7 years or older: give a booster dose every 5 years thereafter, as long as risk remains.
For age 7 through 23 months who have not initiated a series of MenACWY	If age 7–8 months, initiate 2-dose series of Menveo <sup>3</sup> or, if age 9–23 months, give either Menveo or Menactra. <sup>4</sup> Separate the 2 doses by at least 12 weeks.	
For ages 2 years and older	Give 2 doses of MenACWY (any vaccine), 8 weeks apart. <sup>4,7</sup>	
<b>People with HIV infection or functional or anatomic asplenia (including sickle cell disease)</b>		
For age 2 through 6 months	Give 3 doses of Menveo, 8 weeks apart, and a 4th dose at age 12–18 months. If possible, vaccination should begin at age 2 months.	If primary vaccination is completed before the 7th birthday: give one booster dose 3 years after primary series, then every 5 years thereafter. <sup>4</sup>  If primary vaccination is completed at age 7 years or older: give a booster dose every 5 years thereafter.
For age 7 through 23 months who have not initiated a series of MenACWY-CRM	Give 2 doses of Menveo. <sup>3</sup> Separate the 2 doses by at least 12 weeks.	
For ages 2 years and older	Give 2 doses of MenACWY (any vaccine), 8 weeks apart. If using Menactra, give dose #1 at least 4 weeks after final dose of PCV13. <sup>4,7</sup>	

### FOOTNOTES

- If available, use the same vaccine product for all doses in the series given to infants, including the booster doses.
- Seek advice of local public health authorities to determine if vaccination is recommended.
- If initiating vaccination with Menveo in a child age 7 through 23 months, dose 2 should be given no younger than age 12 months.
- If Menactra is to be administered to a child with increased risk for meningococcal disease, it should be given either before, at the same visit, or at least 6 months after DTaP. Menveo and MenQuadfi can be given at any time before or after DTaP.
- If child age 7 through 23 months will enter an endemic area in less than 3 months, give doses as close as 2 months apart.
- Persistent deficiency of complement components C3, C5–C9, properdin, factor D, or factor H caused by an immune system disorder or by taking a complement inhibitor (Soliris [eculizumab] or Ultomiris [ravulizumab]).
- If the person has a history of 1 dose of MenACWY at the time of diagnosis with a high-risk condition for which a 2-dose primary series is recommended, give dose 2, then boost every 5 years as long as risk remains.