

Administering Vaccines to Adults:


Dose, Route, Site, and Needle Size

Vaccine	Dose	Route
COVID-19 For product and dosage information for COVID-19 vaccines, see Immunize.org's "Checklist of Current Versions of U.S. COVID-19 Vaccination Guidance and Clinic Supply Tools" at www.immunize.org/catg.d/p3130.pdf		IM
Hepatitis A (HepA)	≤18 yrs: 0.5 mL; ≥19 yrs: 1.0 mL	IM
Hepatitis B (HepB)	Engerix-B; Recombivax HB ≤19 yrs: 0.5 mL, ≥20 yrs: 1.0 mL Heplisav-B: ≥18 yrs: 0.5 mL	IM
HepA-HepB (Twinrix)	≥18 yrs: 1.0 mL	IM
Human papillomavirus (HPV)	0.5 mL	IM
Influenza, live attenuated (LAIV)	0.2 mL (0.1 mL in each nostril)	Intranasal spray
Influenza, injectable - Egg-based (IIV, Fluarix, FluLaval, Fluzone) - Cell-culture based (ccIV, Flucelvax) - Recombinant (RIV, Flublok) - Adjuvanted (aIIV, Fluad) - High-dose (HD-IIV, Fluzone High Dose)	0.5 mL	IM
	0.38 mL	
Measles, Mumps, Rubella (MMR)	0.5 mL	MMR II (Merck) Priorix (GSK)
		IM or Subcut Subcut
Meningococcal serogroups A, C, W, Y (MenACWY)	0.5 mL	IM
Meningococcal serogroup B (MenB)	0.5 mL	IM
Meningococcal serogroups A, B, C, W, Y (MenABCWY)	0.5 mL	IM
Mpox (Jynneos)	0.5 mL	Subcut ¹
Pneumococcal conjugate (PCV)	0.5 mL	IM
Pneumococcal polysaccharide (PPSV23)	0.5 mL	IM or Subcut
Respiratory Syncytial Virus (RSV) vaccine	0.5 mL	IM
Tetanus, Diphtheria (Td) with Pertussis (Tdap)	0.5 mL	IM
Varicella (VAR)	0.5 mL	IM or Subcut
Zoster (RZV, Shingrix)	0.5 ² mL	IM


1 Administration of Jynneos subcut (0.5 mL) is preferred. If an adult patient desires intradermal administration (0.1 mL) it is permitted under FDA emergency use authorization (see www.fda.gov/media/160774/download).

2 The Shingrix (RZV) vial may contain more than 0.5 mL. Do not administer more than 0.5 mL.


Intramuscular (IM) injection



Subcutaneous (Subcut) injection



Intranasal (NAS) administration of FluMist (LAIV) vaccine



NOTE: Always refer to the package insert (found at www.immunize.org/official-guidance/fda/pkg-inserts) included with each biologic for complete vaccine administration information.

Injection Site and Needle Size

* Alternate needle lengths may be used if the skin is stretched tightly and subcutaneous tissues are not bunched, as follows: a) a 5/8" needle for patients weighing less than 130 lbs (<60 kg) for IM injection in the deltoid muscle only, or b) a 1" needle for IM injection in the antero-lateral thigh muscle for adults of any weight.

Subcutaneous (Subcut) injection – Use a 23–25 gauge, 5/8" needle. Inject in fatty tissue over triceps.		
Intramuscular (IM) injection – Use a 22–25 gauge needle. Choose the needle length and site as indicated below:		
BIOLOGICAL SEX AND WEIGHT OF PATIENT	NEEDLE LENGTH	INJECTION SITE
Female or male less than 130 lbs	5/8" [*] –1"	Deltoid muscle of arm
Female or male 130–152 lbs	1"	
Female 153–200 lbs	1"–1½"	
Male 153–260 lbs		
Female 200+ lbs	1½"	
Male 260+ lbs		
Female or male, any weight	1" [*] –1½"	Anterolateral thigh

