

Haemophilus influenzae type b vaccination**Special situations**

- **Anatomical or functional asplenia (including sickle cell disease):** 1 dose Hib if previously did not receive Hib; if elective splenectomy, 1 dose Hib, preferably at least 14 days before splenectomy
- **Hematopoietic stem cell transplant (HSCT):** 3-dose series Hib 4 weeks apart starting 6–12 months after successful transplant, regardless of Hib vaccination history

Hepatitis A vaccination**Routine vaccination**

- **Not at risk but want protection from hepatitis A** (identification of risk factor not required): 2-dose series HepA (Havrix 6–12 months apart or Vaqta 6–18 months apart [minimum interval: 6 months]) or 3-dose series HepA-HepB (Twinrix at 0, 1, 6 months [minimum intervals: 4 weeks between doses 1 and 2, 5 months between doses 2 and 3])

Special situations

- **At risk for hepatitis A virus infection:** 2-dose series HepA or 3-dose series HepA-HepB as above
 - **Chronic liver disease**
 - **Clotting factor disorders**
 - **Men who have sex with men**
 - **Injection or non-injection drug use**
 - **Homelessness**
 - **Work with hepatitis A virus** in research laboratory or nonhuman primates with hepatitis A virus infection
 - **Travel in countries with high or intermediate endemic hepatitis A**
 - **Close personal contact with international adoptee** (e.g., household, regular babysitting) in first 60 days after arrival from country with high or intermediate endemic hepatitis A (administer dose 1 as soon as adoption is planned, at least 2 weeks before adoptee's arrival)

Hepatitis B vaccination**Routine vaccination**

- **Not at risk but want protection from hepatitis B** (identification of risk factor not required): 2- or 3-dose series HepB (2-dose series Heplisav-B at least 4 weeks apart [2-dose series HepB only applies when 2 doses of Heplisav-B are used at least 4 weeks apart] or 3-dose series Engerix-B or Recombivax HB at 0, 1, 6 months [minimum intervals: 4 weeks between doses 1 and 2, 8 weeks between doses 2 and 3, 16 weeks between doses 1 and 3]) or 3-dose series HepA-HepB (Twinrix at 0, 1, 6 months [minimum intervals: 4 weeks between doses 1 and 2, 5 months between doses 2 and 3])

Special situations

- **At risk for hepatitis B virus infection:** 2-dose (Heplisav-B) or 3-dose (Engerix-B, Recombivax HB) series HepB, or 3-dose series HepA-HepB as above
 - **Hepatitis C virus infection**
 - **Chronic liver disease** (e.g., cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, alanine aminotransferase [ALT] or aspartate aminotransferase [AST] level greater than twice upper limit of normal)
 - **HIV infection**
 - **Sexual exposure risk** (e.g., sex partners of hepatitis B surface antigen (HBsAg)-positive persons; sexually active persons not in mutually monogamous relationships, persons seeking evaluation or treatment for a sexually transmitted infection, men who have sex with men)
 - **Current or recent injection drug use**
 - **Percutaneous or mucosal risk for exposure to blood** (e.g., household contacts of HBsAg-positive persons; residents and staff of facilities for developmentally disabled persons; health care and public safety personnel with reasonably anticipated risk for exposure to blood or blood-contaminated body fluids; hemodialysis, peritoneal dialysis, home dialysis, and predialysis patients; persons with diabetes mellitus age younger than 60 years and, at discretion of treating clinician, those age 60 years or older)
 - **Incarcerated persons**
 - **Travel in countries with high or intermediate endemic hepatitis B**

Human papillomavirus vaccination**Routine vaccination**

- **Females through age 26 years and males through age 21 years:** 2- or 3-dose series HPV vaccine depending on age at initial vaccination; males age 22 through 26 years may be vaccinated based on individual clinical decision (HPV vaccination routinely recommended at age 11–12 years)
- **Age 15 years or older at initial vaccination:** 3-dose series HPV vaccine at 0, 1–2, 6 months (minimum intervals: 4 weeks between doses 1 and 2, 12 weeks between doses 2 and 3, 5 months between doses 1 and 3; repeat dose if administered too soon)
- **Age 9 through 14 years at initial vaccination and received 1 dose, or 2 doses less than 5 months apart:** 1 dose HPV vaccine
- **Age 9 through 14 years at initial vaccination and received 2 doses at least 5 months apart:** HPV vaccination complete, no additional dose needed
- If completed valid vaccination series with any HPV vaccine, no additional doses needed

Special situations

- **Immunocompromising conditions (including HIV infection) through age 26 years:** 3-dose series HPV vaccine at 0, 1–2, 6 months as above
- **Men who have sex with men and transgender persons through age 26 years:** 2- or 3-dose series HPV vaccine depending on age at initial vaccination as above
- **Pregnancy through age 26 years:** HPV vaccination not recommended until after pregnancy; no intervention needed if vaccinated while pregnant; pregnancy testing not needed before vaccination