

Hepatitis A vaccination

(minimum age: 12 months for routine vaccination)

Routine vaccination

- 2-dose series (**Havrix** 6–12 months apart or **Vaqta** 6–18 months apart, minimum interval 6 months); a series begun before the 2nd birthday should be completed even if the child turns 2 before the second dose is administered.

Catch-up vaccination

- Anyone 2 years of age or older may receive HepA vaccine if desired. Minimum interval between doses: 6 months
- Adolescents 18 years and older may receive the combined HepA and HepB vaccine, **Twinrix**, as a 3-dose series (0, 1, and 6 months) or 4-dose series (0, 7, and 21–30 days, followed by a dose at 12 months).

International travel

- Persons traveling to or working in countries with high or intermediate endemic hepatitis A (wwwnc.cdc.gov/travel/):
 - **Infants age 6–11 months:** 1 dose before departure; revaccinate with 2 doses, separated by 6–18 months, between 12 to 23 months of age.
 - **Unvaccinated age 12 months and older:** 1st dose as soon as travel considered

Special situations

At risk for hepatitis A infection: 2-dose series as above

- **Chronic liver disease**
- **Clotting factor disorders**
- **Men who have sex with men**
- **Injection or non-injection drug use**
- **Homelessness**
- **Work with hepatitis A virus** in research laboratory or nonhuman primates with hepatitis A infection
- **Travel** in countries with high or intermediate endemic hepatitis A
- **Close, personal contact with international adoptee** (e.g., household or regular babysitting) in first 60 days after arrival from country with high or intermediate endemic hepatitis A (administer dose 1 as soon as adoption is planned, at least 2 weeks before adoptee's arrival)

Hepatitis B vaccination

(minimum age: birth)

Birth dose (monovalent HepB vaccine only)

- **Mother is HBsAg-negative:** 1 dose within 24 hours of birth for **all** medically stable infants $\geq 2,000$ grams. Infants $< 2,000$ grams: administer 1 dose at chronological age 1 month or hospital discharge.

• **Mother is HBsAg-positive:**

- Administer **HepB vaccine** and **0.5 mL of hepatitis B immune globulin (HBIG)** (at separate anatomic sites) within 12 hours of birth, regardless of birth weight. For infants $< 2,000$ grams, administer 3 additional doses of vaccine (total of 4 doses) beginning at age 1 month.
- Test for HBsAg and anti-HBs at age 9–12 months. If HepB series is delayed, test 1–2 months after final dose.

• **Mother's HBsAg status is unknown:**

- Administer **HepB vaccine** within 12 hours of birth, regardless of birth weight.
- For infants $< 2,000$ grams, administer **0.5 mL of HBIG** in addition to HepB vaccine within 12 hours of birth. Administer 3 additional doses of vaccine (total of 4 doses) beginning at age 1 month.
- Determine mother's HBsAg status as soon as possible. If mother is HBsAg-positive, administer **0.5 mL of HBIG** to infants $\geq 2,000$ grams as soon as possible, but no later than 7 days of age.

Routine series

- 3-dose series at 0, 1–2, 6–18 months (use monovalent HepB vaccine for doses administered before age 6 weeks)
- Infants who did not receive a birth dose should begin the series as soon as feasible (see Table 2).
- Administration of **4 doses** is permitted when a combination vaccine containing HepB is used after the birth dose.
- **Minimum age** for the final (3rd or 4th) dose: 24 weeks
- **Minimum intervals:** dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 8 weeks / dose 1 to dose 3: 16 weeks (when 4 doses are administered, substitute "dose 4" for "dose 3" in these calculations)

Catch-up vaccination

- Unvaccinated persons should complete a 3-dose series at 0, 1–2, 6 months.
- Adolescents age 11–15 years may use an alternative 2-dose schedule with at least 4 months between doses (adult formulation **Recombivax HB** only).
- Adolescents 18 years and older may receive a 2-dose series of HepB (**Heplisav-B**) at least 4 weeks apart.
- Adolescents 18 years and older may receive the combined HepA and HepB vaccine, **Twinrix**, as a 3-dose series (0, 1, and 6 months) or 4-dose series (0, 7, and 21–30 days, followed by a dose at 12 months).
- For other catch-up guidance, see Table 2.

Human papillomavirus vaccination

(minimum age: 9 years)

Routine and catch-up vaccination

- HPV vaccination routinely recommended for all adolescents **age 11–12 years (can start at age 9 years)** and through age 18 years if not previously adequately vaccinated
- 2- or 3-dose series depending on age at initial vaccination:
 - **Age 9 through 14 years at initial vaccination:** 2-dose series at 0, 6–12 months (minimum interval: 5 months; repeat dose if administered too soon)
 - **Age 15 years or older at initial vaccination:** 3-dose series at 0, 1–2 months, 6 months (minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 12 weeks / dose 1 to dose 3: 5 months; repeat dose if administered too soon)
- If completed valid vaccination series with any HPV vaccine, no additional doses needed

Special situations

- **Immunocompromising conditions, including HIV infection:** 3-dose series as above
- **History of sexual abuse or assault:** Start at age 9 years
- **Pregnancy:** HPV vaccination not recommended until after pregnancy; no intervention needed if vaccinated while pregnant; pregnancy testing not needed before vaccination

Inactivated poliovirus vaccination

(minimum age: 6 weeks)

Routine vaccination

- 4-dose series at ages 2, 4, 6–18 months, 4–6 years; administer the final dose on or after the 4th birthday and at least 6 months after the previous dose.
- 4 or more doses of IPV can be administered before the 4th birthday when a combination vaccine containing IPV is used. However, a dose is still recommended after the 4th birthday and at least 6 months after the previous dose.

Catch-up vaccination

- In the first 6 months of life, use minimum ages and intervals only for travel to a polio-endemic region or during an outbreak.
- IPV is not routinely recommended for U.S. residents 18 years and older.

Series containing oral polio vaccine (OPV), either mixed OPV-IPV or OPV-only series:

- Total number of doses needed to complete the series is the same as that recommended for the U.S. IPV schedule. See www.cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm?s_cid=mm6601a6_w.